

AD \_\_\_\_\_

COOPERATIVE AGREEMENT NUMBER DAMD17-95-2-5003

TITLE: Collaborative Research and Support of Fitzsimmons Army  
Medical Center Defense Women's Health Research Program Projects

SUBTITLE: The Effect of Region-Specific Resistance Exercises on  
Bone Mass in Premenopausal Military Women (Protocol #8)

PRINCIPAL INVESTIGATION: Mulligan, Hugh L.  
McDermott, Michael T., COL

CONTRACTING ORGANIZATION: Facilitators of Applied Clinical Trials  
San Antonio, Texas 78216

REPORT DATE: 30 June 1996

TYPE OF REPORT: Final

PREPARED FOR: Commander  
U.S. Army Medical Research and Materiel Command  
Fort Detrick, Frederick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;  
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

19970908 051

**Form Approved**  
**OMB No. 0704-0188**

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 30 June 1996	3. REPORT TYPE AND DATES COVERED Final (1 Feb 95 - 30 Jun 96)	
4. TITLE AND SUBTITLE Collaborative Research and Support of Fitzsimmons Army Medical Center Defense Women's Health Research Program Projects SUBTITLE: The Effect of Region-Specific Resistance Exercises on Bone Mass ...		5. FUNDING NUMBERS  DAMD17-95-2-5003	
6. AUTHOR(S)  Mulligan, Hugh L. McDermott, Michael T., COL		8. PERFORMING ORGANIZATION REPORT NUMBER  Protocol #8	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Facilitators of Applied Clinical Trials San Antonio, Texas 78216			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Commander U.S. Army Medical Research and Materiel Command Fort Detrick, Frederick, MD 21702-5012		10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES			
12a. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for public release; distribution unlimited		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200)  The attainment of peak bone mass during the premenopausal years is critical in preventing osteoporosis later in life. The purpose of this study was to determine if peak bone mass can be improved after age 20, the age near which peak bone mass is usually reached, and to compare the effects of region specific resistance exercise with aerobic exercise. Forty three volunteers completed six months of the protocol. Thirty individuals completed twelve months with fifteen in each group. Bone density was shown to significantly increase in the spine, femoral neck and radius by 12 months in both groups. The resistance exercise group had significantly higher densities at the femoral neck and mid-radius at 12 months.  DTIC QUALITY INSPECTED 4			
14. SUBJECT TERMS Nautilus, aerobic, exercise, osteoporosis, bone mass, premenopausal, calcium  Defense Women's Health Research Program		15. NUMBER OF PAGES 13 16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited

## FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the US Army.

Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

~~MTM~~ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

*Michael T. McDermott*  
PI - Signature                      Date

25 June 1996

## TABLE OF CONTENTS

Cover.....	i
Report Documentation Page.....	ii
Foreword.....	iii
Table of Contents.....	1
Introduction.....	2
Body.....	4
Conclusions.....	6
References.....	7
Tables.....	8

## Introduction

Postmenopausal osteoporosis is a growing health concern in the United States ultimately generating over \$10 billion annually in health care costs. Development of osteoporosis in later life is known to reflect lack of bone health in early life. Peak bone mass is achieved between ages 18 and 35 and becomes the reserve upon which further aging and menopause will draw over the remainder of an individual's life.

Prevention of postmenopausal osteoporosis is recognized to be dependent on calcium intake, physical activity and genetics, as these factors determine the maximum peak bone mass in young women. Skeletal effects of exercise have been the subject of extensive recent literature reviews<sup>(1-3)</sup>. Cross-sectional studies have suggested higher bone mass developing in regions of specific bone loading exercises. However, these studies were flawed by self selection and many inconsistencies which have rendered the overall results inconclusive. Prospective longitudinal studies have shown benefit of vigorous aerobic and resistance exercises, especially in the loaded areas. These benefits appear to be true for both premenopausal and postmenopausal women. These studies were flawed, however, by inadequate randomization. One cross-sectional study of premenopausal women showed weight training to be more effective than aerobic<sup>(4)</sup>. However, four well designed prospective studies on pre- and postmenopausal women have shown varying results which include an increase<sup>(5)</sup>, minimal increase<sup>(6)</sup>, no

change<sup>(7)</sup>, and a decrease<sup>(8)</sup> of bone mass in the loaded regions.

In this study 82 premenopausal women ages 18-40 were enrolled in a one year period in either a resistance-exercise (Nautilus) group or an aerobic group.

Individuals enrolled in the Nautilus group performed exercises specifically designed to strengthen the lower back, abdomen, hips and arms in three 30-minute sessions per week. All individuals in the study completed a questionnaire on calcium intake and were given calcium supplements or were asked to increase dietary calcium intake to fulfill current recommendations for calcium intake of 1000 mg/day in premenopausal women. Bone density determinations were done on entry into the study and at six months and at one year for the lumbar spine, hip and radius which were the areas most likely to be affected by osteoporotic fractures.

## Body

Eighty-two premenopausal women ages 18-40 were initially enrolled in the study and calcium intake and baseline laboratory data collected. The individuals were assigned to the aerobic exercise or Nautilus study groups (according to individual preference for the type of exercise being done), with 44 entering the Nautilus group and 38 entering in the aerobic group. Baseline demographic data on the two groups is provided in Table 1. All individuals in the study were instructed on their prescribed exercises and were given log sheets to record the exercises done, with individual notation for each session including time of session, weight lifted, repetitions and weekly frequency of exercise sessions. Baseline bone density determinations were obtained on 76 subjects and 6 month determinations on 43 subjects. Thirty subjects completed 12 month determinations, with 15 individuals in each group. Six subjects did not complete bone densities and were lost early in the study. Exercise logs were reviewed at the three month interval visits. Also an assessment of calcium intake was completed at these visits and compliance with dietary and supplemental calcium recommendations noted.

An additional parameter was added to obtain bioelectric impedance testing of body fat content on all individuals. This addition was approved through appropriate channels and the measurements were obtained after the participants had been properly counseled and signed a separate consent form.

Reasons for dropouts during the study include: pregnancy (2), medical problems (7), retirement from the Army (3), movement to new assignment (14), lack of time (19), temporary duty away (1).

Delays in the protocol approval process resulted in a six month delay in starting this study and therefore an application for an extension of the study period was approved.

Changes in bone density were calculated using paired t-tests for measurements at zero and six months and zero and twelve months. Statistically significant increases in bone density were found in both groups at all three sites. See Table 2.

The groups were then compared with a z-test for means to compare percent changes in bone density. See Table 3.



## Conclusions

Thirty premenopausal women completed a one year trial comparing aerobic to resistance-exercise for influence on bone density. Both groups showed significant increases in bone density compared to baseline at all 3 sites after both 6 months and 12 months of exercise. The two groups were compared by assessing percent changes in bone density from a baseline at six and twelve months. At six months bone density had increased significantly more at L2-4 in the resistance exercise group but more at the mid-radius in the aerobic group. At twelve months the resistance group showed higher bone densities at the femoral neck and mid-radius but the two groups were not significantly different at L2-4. Site specific analysis of resistance exercises may further delineate the benefits of specific resistance exercises.

Our data suggest that both aerobic and resistance exercises improve bone density in premenopausal women and further, that a moderate program of resistance exercise which emphasizes the hip girdle and arm regions has a greater effect at the mid-radius and femoral neck than does aerobic exercise. Recommending resistance exercises to premenopausal women, by increasing peak premenopausal bone mass, may afford increased protection against hip and wrist fractures in the postmenopausal years.

## Reference

1. Sinaki M. Exercise and osteoporosis. Arch Phys Med Rehabil 70:220-9, 1989.
2. Gutin B, Kasper MJ. Can vigorous exercise play a role in osteoporosis prevention? a review. Osteoporosis Int 2:55-69, 1992.
3. Drinkwater BL. Exercise in the prevention of osteoporosis. Osteoporosis Int 1:S169-S171, 1993.
4. Heinrich CH, Going SB, Pamenter RW, Perry CD, Boyden TW, Lohman TG. Bone mineral content of cyclically menstruating female resistance and endurance trained athletes. Med Sci Sports Exerc 22:558-63, 1990.
5. Snow-Harter C, Bouxsein ML, Lewis BT, Carter DR, Marcus R. Effects of resistance and endurance exercise on bone mineral status of young women: a randomized exercise intervention trial. J Bone Min Res 7:761-9, 1992.
6. Gleeson PB, Protas EJ, LeBlanc AD, Schneider VS, Evans HJ. Effects of weight lifting on bone mineral density in premenopausal women. J Bone Min Res 5:153-8, 1990.
7. Peterson SE, Peterson MD, Raymond G, Gilligan C, Checovich MM, Smith EL. Muscular strength and bone density with weight training in middle-aged women. Med Sci Sports Exerc 23:499-504, 1991.
8. Rockwell JC, Sorensen AM, Baker S, et al. Weight training decreases vertebral bone density in premenopausal women: a prospective study. J Clin Endocrinol Metab 71:988-993, 1990.

## EXERCISE.XLS

## EXERCISE STUDY

TABLE 1 DEMOGRAPHIC DATA

		AEROBIC (A)		NAUTILUS (N)	
		Mean	Range	SE	n
Age	(A)	31.6	21-40	0.92	38
	(N)	31.8	19-40	0.75	44
Height	(A)	64.7	56.8-69.8	0.41	38
	(N)	65.3	60.5-74.3	0.39	44
Weight	(A)	144.1	112-236	4.1	38
	(N)	143.3	100-183	3.15	44
% body fat	(A)	28.1	14-45	1.2	37
	(N)	27.1	15-40	1.1	42
Scapular skin fold	(A)	16.2		0.98	38
	(N)	14.3		0.65	44
Ca intake	(A)	835	173-2521	98.7	38
	mg/d (N)	780	144-2161	66.2	44
Bone density mg/cm <sup>2</sup>					
L2-4	(A)	1.218	1.063-1.429	0.017	33
	(N)	1.238	0.946-1.586	0.025	41
Fem. neck	(A)	1.047	0.797-1.339	0.023	34
	(N)	1.062	0.830-1.463	0.02	42
Mid-radius	(A)	0.707	0.632-0.870	0.0086	34
	(N)	0.71	0.606-0.824	0.0073	42
Race %		Caucasian	Black	Hispanic	Other
	(A)	65.8	15.8	10.5	7.9
	(N)	89.5	15.8	7.9	2.6

## EXERCISE STUDY

TABLE 2 BONE MINERAL DENSITIES

AEROBIC  
(n = 22)

	Initial	6 Months	p
L2-4	1.223	1.235	0.06
Fem. neck	1.047	1.099	<0.001
Mid-radius	0.709	0.755	<0.001

NAUTILUS  
(n = 21)

L2-4	1.181	1.206	0.003
Fem. neck	1.028	1.082	<0.001
Mid-radius	0.707	0.749	<0.001

AEROBIC  
(n = 15)

	Initial	12 Months	p
L2-4	1.232	1.254	0.003
Fem. neck	1.036	1.064	0.03
Mid-radius	0.714	0.762	<0.001

NAUTILUS  
(n = 15)

L2-4	1.184	1.21	0.04
Fem. neck	1.01	1.06	0.03
Mid-radius	0.705	0.76	<0.001

## EXERCISE STUDY

TABLE 3      PERCENT CHANGE FROM BASELINE  
IN BONE DENSITIES

6 MONTHS		Aerobic	Nautilus	p
	L2-4	0.9	1.93	<0.001
	Fem. neck	5.16	5.46	0.16
	Mid-radius	6.62	5.97	0.02
12 MONTHS				
	L2-4	1.71	2.16	0.11
	Fem. neck	2.76	5.23	<0.001
	Mid-radius	7.03	7.71	0.03